arXiv:1207.6810v2 [gr-qc] 15 Aug 2012

Splitting singular surfaces

Marcos A. Ramirez

Instituto de Física Enrique Gaviola, FaMAF,
Universidad Nacional de Córdoba, (5000) Córdoba, Argentina

In this paper we first consider a singular timelike spherical hypersurface embedded in a D-dimensional spherically symmetric bulk space-time. We analyze the different possibilities regarding the orientation of the gradient of the standard r coordinate in relation to the shell. The matter content of the shell is described by a number of non-interacting spherically symmetric matter fields. We analyze the dynamics according to Einstein's equations for matter satisfying certain energy conditions. Then we show that, under some circumstances, the system at a given proper time becomes unstable against an infinitesimal separation of non-interacting constituents. In particular, we consider the case where the shell is filled with collisionless particles that move in trajectories of constant angular momentum, and show that, for shells whose angular momentum distribution is discrete, the instability that we described before takes place. We also find this instability when the components are fluids with certain equations of state. We give explicit examples and construct solutions that represent a shell that splits into two shells. Then we extend those results for 5-dimensional Schwarzschild-AdS bulk space-times, which is a typical scenario for brane-world models, and show that the same kind of stability analysis and splitting solution can be constructed. We find that a widely proposed family of brane-world models are extremely unstable in this sense. Finally, we discuss possible interpretations of these features and their relation to the initial value problem with concentrated sources.

PACS numbers:

I. INTRODUCTION

Singular surfaces proved to be a very useful tool in constructing toy-models to gain insight into general theoretical problems in general relativity, like the cosmic censorship conjecture [5]. They are also useful in constructing models for a number of concrete astrophysical and cosmological situations ([6], [7]) where shock-like surfaces formed, or where, as a valid simplifying assumption, some part of the system can be thought of as a singular surface. In this way, we consider that an analysis of general mathematical properties of this kind of objects is important and may have possible relevant astrophysical and cosmological implications. In particular, as a far as we know, an analysis on the non-uniqueness of the evolution of initial data associated with singular hypersurfaces has not been performed.

Inspired by string theory, certain cosmological models that consider the visible universe as a part of a singular 4-dimensional surface embedded in a higher-dimensional space-time appeared [2],[3]. Besides being a theoretical possibility that is worth exploring, brane-world models have been proved to have interesting properties, such a Friedmann-like evolution equations that exhibit an inflationary stage [4]. They represent phenomenological models that can account for the "invisibility" of extra dimensions, and constitute a framework where the holographic principle can be naturally understood.

Outline

We begin with a description of a general shell in a spherically symmetric bulk space-time of an arbitrary number of dimensions in Section II. In Section III we analyze some aspects of the dynamics of these shells, provided their matter-energy content satisfy the dominant energy condition. In Section we develop the stability analysis against separation of non-interacting constituents and find a stability condition for general constituents, and then construct some examples of splitting solutions in Section V. In Section VI, we analyze a splitting scenario that preserves Z_2 symmetry in which a three component shell (with two of its constituents being identical) splits into three shells. Then in Section VII we extend both kind of instabilities to non-Z_2-symmetric and Z_2-symmetric brane-world models respectively. Finally, in Section VIII we comment on possible interpretations of these features in the context of the general initial value problem with concentrated sources. We formulate the following conjecture: *unstable hypersurfaces are not thin-shell limits of thick configurations.*

II. SPHERICALLY SYMMETRIC THIN SHELLS IN $D = n + 2$ DIMENSIONS

We begin with a description of a singular shell embedded in a spherically symmetric vacuum space-time of an arbitrary number of dimensions. The idea of spherical symmetry is originally defined for $n + 1$-dimensional riemannian manifolds: there is a group of isometries isomorphic to the $SO(n)$ group. For lorentzian manifolds, we say that there exists a spacelike foliation such that every riemannian slice is spherically symmetric and the lapse and shift are invariant against the action of the group. Charting the space of group orbits with orthogonal coordinates (x_0, x_1), which is always possible, we may write

$$ds^2 = -f(x_0, x_1)dx_0^2 + h(x_0, x_1)dx_1^2 + r(x_0, x_1)^2 d\Omega_n^2. \tag{1}$$

We have a singular timelike orientable hypersurface Σ embedded in the bulk space-time. Because of the symmetry, the surface should be described by an equation $\Sigma(x^0, x^1) = 0$. In particular, in gaussian coordinates, in a neighborhood of Σ the metric reads,

$$ds^2 = -f(\tau, \eta)d\tau^2 + d\eta^2 + r(\tau, \eta)^2 d\Omega_n^2 \tag{2}$$

where $\eta = 0$ characterizes the surface, and $f(\tau, 0) = 1$ (τ is the shell proper time).

Einstein equations imply junction conditions on the surface [1] that relate the jump of the surface extrinsic curvature with the effective mass-energy tensor on the shell. These are the so-called Israel-Darmois junction conditions. The extrinsic curvature in gaussian coordinates reads,

$$K^i_j = diag\left[\frac{1}{2}\frac{\partial f}{\partial \eta}|_{\eta=0}, \frac{1}{r}\frac{\partial r}{\partial \eta}|_{\eta=0}, .., \frac{1}{r}\frac{\partial r}{\partial \eta}|_{\eta=0}\right] \tag{3}$$

where, and from now on, latin indexes represent coordinates $(\tau, \theta_1, .., \theta_n)$ on the shell. In these coordinates the intrinsic metric takes the form,

$$ds^2_\Sigma = -d\tau^2 + R(\tau)^2 d\Omega_n^2 \tag{4}$$

where $R(\tau) \equiv r(\tau, 0)$.

By virtue of Birkhoff theorem, vacuum solutions of Einstein equations can always be written in the form,

$$ds^2 = -F(r)dt^2 + F(r)^{-1}dr^2 + r^2 d\Omega_n^2 \tag{5}$$

where $F(r) = 1 - 2M/r^{n-1}$, and M is a parameter equivalent to the Misner-Sharp mass inside the *innermost* group orbit. From now on, we assume $R(\tau)^{n-1} > 2M$, unless otherwise stated, since we are not interested in collapsing situations. With these expressions, we can write the extrinsic curvature on a given *side* of the shell in terms of $R(\tau)$ and the parameter M that characterizes the bulk space-time there

$$K^i_j = sign\left(\frac{\partial r}{\partial \eta}|_{\eta=0}\right) diag\left[\frac{F'(R)+2\ddot{R}}{2\sqrt{\dot{R}^2+F(R)}}, \frac{\sqrt{\dot{R}^2+F(R)}}{R}, .., \frac{\sqrt{\dot{R}^2+F(R)}}{R}\right]. \tag{6}$$

In this way, giving M and specifying whether r increases or decreases with η, we get an expression for the extrinsic curvature in terms of the *intrinsic* function $R(\tau)$.

On the other hand, the matter content of the shell is typically described by a tensor S^i_j defined on Σ such that we can formally write the D-dimensional stress-energy tensor as,

$$T^a_b = \delta(\Sigma) S^a_b \tag{7}$$

where the S tensor in shell coordinates takes the form

$$S^i_j = diag[-\rho(\tau), p(\tau), ..., p(\tau)]. \tag{8}$$

So, as a result of the symmetry imposed, we can describe the matter content of the shell as if it were composed of a n-dimensional perfect fluid, whose flow lines follow the trajectories of the comoving observers. Explicitly,

$$S^{ij} = ph^{ij} + (\rho + p)u^i u^j \tag{9}$$

where h_{ij} is the intrinsic metric defined in (4), and $u^i = (\partial/\partial\tau)^i$. If there is not hysteresis, we should be able to write ρ and p as functions of R. In that case, conservation of the source would read,

$$\frac{d\rho}{dR} + \frac{n(\rho+p)}{R} = 0 \tag{10}$$

This equation together with an equation of state $f(\rho, p) = 0$, provided it exists, determine $\rho(R)$ and $p(R)$.

Now we can see that Israel-Darmois junction conditions should relate $R(\tau)$, it first two derivatives, and the parameters M with the matter functions $\rho(R)$ and $p(R)$. Concentrated matter on the shell implies the *discontinuity* of the extrinsic curvature. Looking at (6), we realize that the jump should be ascribed to a difference between the mass parameters for the empty regions at both sides of the shell, which we call M_I and M_{II}, and, eventually, to different signs for $\partial r/\partial\eta$ at both sides. In this paper we consider situations where at least one of the two Schwarzschild regions, say region I, is *interior*, that is, it can be described by an inequality $r < R(t)$ in terms of the standard coordinates for that region. The reason for this choice, which only precludes the case where both regions are *exterior*, will become clear when we analyze the junction conditions. From now on, we choose the η coordinate to decrease when going into region I, which implies $\partial r/\partial\eta|_{\eta=0^-} > 0$. The junction conditions then read,

$$\frac{n}{R}\left(\pm\sqrt{\dot{R}^2+F_{II}} - \sqrt{\dot{R}^2+F_I}\right) = -\kappa\rho \tag{11}$$

$$\pm\frac{F'_{II}+2\ddot{R}}{2\sqrt{\dot{R}^2+F_{II}}} - \frac{F'_I+2\ddot{R}}{2\sqrt{\dot{R}^2+F_I}} + \frac{n-1}{R}\left(\pm\sqrt{\dot{R}^2+F_{II}} - \sqrt{\dot{R}^2+F_I}\right) = \kappa p \tag{12}$$

where $F_i = 1 - 2M_i/r^{n-1}$ and $\pm = sign(\partial r/\partial\eta|_{\eta=0^+})$. The situation not considered by these expressions, the case where I and II are both *exterior* regions, is not generally regarded as *physical* because it would imply a *negative energy density* ρ. In the context of brane cosmology, however, a *brane tension* is assumed, which implies an effective constant energy density on the shell, that could in principle be negative, because it is a non-dynamical entity. But, if that were the case, we would not recover standard cosmology as a low energy (at least since recombination) limit. So even in those speculative scenarios, a negative energy density type of matter-energy is hard to justify phenomenologically. In this way, we will only consider matter that satisfies the dominant energy condition, which is a prescription that includes an eventual positive brane tension. In particular, in the case of a *exterior* region II, a positive effective energy density ρ would imply $M_{II} > M_I$.

From (11) we can obtain an equation of motion that results independent of the sign $\pm$ and reads

$$\frac{1}{2}\dot{R}^2 + V(R) = 0 \quad , \quad V(R) = \frac{1}{2} - \frac{M_I + M_{II}}{2R^{n-1}} - \frac{n^2(M_{II} - M_I)^2}{2\kappa^2\rho^2R^{2n}} - \frac{\kappa^2\rho^2R^2}{8n^2}. \tag{13}$$

But, in obtainig this equation, we have squared some quantities, so it may contain spurious solutions, that is, there could be solutions of (13) which are not solutions of (11) with a definite sign for $\partial r/\partial\eta|_{\eta=0^+}$. Nevertheless, it can be shown that every solution of (13) is always a solution of one of the two versions of (11) (with a specific choice for $\pm$). Therefore, we can construct solutions of the junction conditions by solving (13) and specifying *a posteriori* $\partial r/\partial\eta|_{\eta=0^+}$ accordingly[1]. For a given solution of the equation of motion, $R(\tau)$, the specification that we should made is the following

$$sign(\partial r/\partial\eta|_{\eta=0^+}) = sign(n^2(F_I - F_{II}) - \kappa^2\rho^2R^2) = sign(2n^2(M_{II} - M_I) - \kappa^2\rho^2R^{n+1}). \tag{14}$$

It can be shown that this specification is consistent, because the roots of $2n^2(M_{II} - M_I) - \kappa^2\rho(R)^2R^{n+1}$ are always in *prohibited regions* ($V(R) > 0$) or inside an event horizon $R^{n-1} < 2M_{(I,II)}$. In this way, for a given initial configuration of the shell (for an arbitrary choice of parameters $(R_0, \rho(R_0), M_{II}, M_I)$), we can tell whether region II should be *interior*, or *exterior*, in order to satisfy the junction conditions, and that specification turns out to be consistent throughout the evolution [2]. Now it is clear that giving a function $\rho(R)$ (something that we can find for example imposing an equation of state) and the parameters (M_I,M_{II}), we have a complete description of the possible motions of the shell by solving an equation of motion for a single particle moving in a given potential.

On the other hand, equation (12) implies,

$$\ddot{R} = -\frac{n-1}{2R}\left[1 + \dot{R}^2 \mp \left(\frac{2n}{n-1}\alpha + 1\right)\sqrt{1 + \dot{R}^2 - \frac{2M_I}{R^{n-1}}}\sqrt{1 + \dot{R}^2 - \frac{2M_{II}}{R^{n-1}}}\right] \tag{15}$$

where $\alpha \equiv p/\rho$. This expression will be useful later on.

III. EQUATIONS OF MOTION AND MATTER MODELS

In this Section we analyze general properties of the dynamics of the shell geometry, provided the matter content satisfies an energy condition. In particular, we analyze the dynamics for some matter-energy models of interest: fluids with equation of state $p = \omega\rho$ and Vlasov matter.

---

[1] The general problem of gluing two spherically symmetric space-times by means of an hypersurface was considered in [12].

[2] Inside an event horizon, the r coordinate becomes timelike, space-time is not static, and the signature of $\partial r/\partial\eta$ may change without contradiction with Birkhoff's theorem

Looking at the effective potential (13), we see that the permitted regions for the function $R(\tau)$ are those in which $V(R) < 0$. The structure of these regions depend on the behavior of the function $\rho(R)$, so we have to specify the matter model and its parameters in order to describe possible motions. The condition $V(R) < 0$ can be explicitly written,

$$4n^2\kappa^2\rho^2R^{2n} - 4n^2\kappa^2(M_I + M_{II})\rho^2R^{n+1} - 4n^4(M_{II} - M_I)^2 - \kappa^4\rho^4R^{2(n+1)} < 0 \tag{16}$$

where it is clear that for R small enough, this inequality is always satisfied, so there are always "collapsing" solutions for the equation of motion (13) provided $R(\tau_0)$ is small enough.

The asymptotic behavior of the potential when $R \to \infty$ depends on the matter model as follows. If we suppose that matter satisfies *the dominant energy condition*, that is $\rho \geq |p|$, then equation (29) implies

$$-2n \leq \frac{d(ln(\rho))}{d(ln(R))} \leq 0. \tag{17}$$

So, giving an initial value $\rho_0 \equiv \rho(R_0)$, we have for $R > R_0$

$$\rho_0\left(\frac{R_0}{R}\right)^{2n} \leq \rho(R) \leq \rho_0. \tag{18}$$

Considering the function $\alpha(R) \equiv p(R)/\rho(R)$, the dominant energy condition then implies $-1 \leq \alpha(R) \leq 1$. We assume $n \geq 2$ and that $\alpha(R)$ has a limit α_∞ when R goes to infinity, which implies that ρ acquires the asymptotic form $\rho \approx CR^{-n(1+\alpha_\infty)}$. According to (14), we can see that if $\alpha_\infty > -(n-1)/2n$, for R large enough, region II would be *exterior*, and if $\alpha_\infty < -(n-1)/2n$ then it would be *interior.* In the particular case $\alpha_\infty = -(n-1)/2n$, region II character would be determined by the sign of $2n^2(M_{II} - M_I) - \kappa^2C^2$. The asymptotic behavior of $V(R)$ in the limit $R \to \infty$ can be classified as follows (in all these cases A stands for different constants that depend on the parameters).

- $-1 \leq \alpha_\infty < -(n-1)/n$

  Then $V(R) \approx -\frac{\kappa^2\rho^2R^2}{8n^2} \to -AR^{2(1-n(1+\alpha_\infty))}$. In this case $R(\tau)$ is unbounded, provided $R(\tau_0)$ is large enough.

- $\alpha_\infty = -(n-1)/n$

  Then $V(R) \approx \frac{4n^2-\kappa^2\rho^2R^2}{8n^2} \to cte$. The condition for having an unbounded $R(\tau)$ solution is an inequality involving the model parameters.

- $-(n-1)/n < \alpha_\infty \leq 0$

  Then $V(R) \to \frac{1}{2}$. There is always an upper bound for $R(\tau)$.

- $\alpha_\infty = 0$

  Then $V(R) \approx \frac{\kappa^2\rho^2R^{2n}-n^2(M_{II}-M_I)^2}{2\kappa^2\rho^2R^{2n}} \to cte$. Once again the condition for having an unbounded $R(\tau)$ is an inequality involving the model parameters.

- $0 < \alpha_\infty \leq 1$

  Then $V(R) \approx -\frac{n^2(M_{II}-M_I)^2}{2\kappa^2\rho^2R^{2n}} \to -AR^{2n\alpha_\infty}$. There are always unbounded solutions.

Now it is clear that not for every matter model there exist unbounded solutions, so the limit $R \to \infty$ not always makes sense. Anyway, this limit, provided it exists, is important because of the fact that for large R we may find Newtonian or low-energy limits for the dynamics of the shell. For example in the context of brane-world cosmologies we hope to recover the standard Friedmann equations for R large in some sense. On the other hand, the existence of unbounded solutions will be useful to construct "unstable-against-separation" solutions, as we will see in the next Sections.

It will also be useful to stress the fact that if we have k non-interacting matter fields (so we can write $\rho = \sum_{i=1}^{k} \rho_i$ and $p = \sum_{i=1}^{k} p_i$), each with its own conservation equation and its corresponding $\alpha_{i\infty}$, then $\alpha_\infty = min\{\alpha_{i\infty}\}$. In the next subsections we analyze some characteristics of $V(R)$ for specific matter models.

1. ***Fluid with equation of state*** $p = \omega\rho$

Equation (29) implies for this family of matter models

$$\rho(R) = \rho_0 \left(\frac{R_0}{R}\right)^{n(1+\omega)} . \tag{19}$$

In this case $V(R)$ can be written

$$V(R) = \frac{1}{2} - \frac{M_I + M_{II}}{2R^{n-1}} - \frac{n^2(M_{II} - M_I)R^{2n\omega}}{2C^2} - \frac{C^2}{8n^2 R^{2(n(1+\omega)-1)}} \tag{20}$$

where $C \equiv \kappa\rho_0 R_0^{n(1+\omega)}$. The dominant energy condition then implies $-1 \leq \omega \leq 1$, and we have $\alpha(R) = \alpha_\infty = \omega$. The character of region II should be determined by the sign of the expression

$$2n^2(M_{II} - M_I)R^{n(1+2\omega)-1} - \kappa^2 C^2. \tag{21}$$

We can see that this expression is a monotonically increasing function of R for $\omega > -(n-1)/2n$, a constant for $\omega = -(n-1)/2n$, and a monotonically decreasing function for $\omega < -(n-1)/2n$. In this way, for large R, in the first case region II should be exterior, and interior in the latter case. We now comment on some specific and relevant cases.

- $\omega = -1$ represents a "cosmological constant fluid" or brane tension. We have $\rho = \rho_0$, and there are always unbounded $R(\tau)$ solutions. For these unbounded solutions, region II must be an *interior* Schwarzschild region. For certain parameters $(n, M_I, M_{II}, \kappa C)$ there are no prohibited regions.

- $\omega = 0$ represents dust, may be the simplest matter model of all. We have $\rho = CR^{-n}$, and the condition for having unbounded $R(\tau)$ is now $M_{II} - M_I \geq C^2/n^2$, otherwise we would have an upper bound. Region II must be an *exterior* Schwarzschild region for the unbounded solutions. $V(R)$ results a monotonically increasing function, so there could be at most one turning point, and if the parameters allow unbounded solutions, then there are no prohibited regions.

- $\omega = 1/n$ represents a radiation shell. We have $\rho = CR^{-(n+1)}$, and there are always unbounded solutions. For these unbounded solutions, region II must be *exterior.* Also in this case, for certain parameters there are no prohibited regions.

2. ***Vlasov matter***

Another simple and relevant matter model used for many astrophysical situations is Vlasov matter [15]. It has the advantage that its range of applicability is phenomenologically clear: a gravitating system can be thought of as an Einstein-Vlasov system whenever it is composed of many constituents with low collision probability. It is defined as an ensemble of autogravitating collisionless particles that interact with each other only by means of the curvature that they generate as a whole. In particular, they follow geodesic trajectories. In this context, we have to specify what do we mean with "geodesic" because of the discontinuity of the metric connection on the shell. Furthermore, the curves contained within the shell can not be in general geodesics of neither connection (see [16] and [17] for a discussion in the context of brane cosmology). The alternative we have is to impose that *the particles follow geodesics of the induced metric*, something that sounds reasonable, specially taking into account that in this symmetry such geodesics are simply the trajectories of conserved angular momentum within the shell. The system we are considering in this subsection is a D-dimensional generalization of the four-dimensional case analyzed in [9].

If all particles are identical, the S tensor and the particle density current can be written,

$$S^{ij} = -\mu \int f(x,p)\sqrt{-h}u^i u^j \frac{dp^1..dp^n}{p_0} \quad , \quad N^i = \int f(x,p)\sqrt{-h}u^i \frac{dp^1..dp^n}{p_0} \tag{22}$$

where $f(x,p)$ is the distribution of the number of particles in the tangent bundle and μ is the particles proper mass. Considering the symmetry and the conservation of the distribution of the number of particles with angular momentum modulus L (which we call $n(L)$), it can be shown that the independent components of the S tensor take the form,

$$\rho(R) = \frac{\mu}{S_n R^{n+1}} \int n(L)\sqrt{R^2+L^2}dL \quad , \quad p(R) = \frac{\mu}{n S_n R^{n+1}} \int \frac{n(L)L^2}{\sqrt{R^2+L^2}}dL \tag{23}$$

where S_n is the surface of a n-sphere of unit radius.

Now we define,

$$f(R) \equiv \int n(L)\sqrt{R^2+L^2}dL \tag{24}$$

so the equation of motion for this shell can be written,

$$V(R) = \frac{1}{2} - \frac{M_I + M_{II}}{2R^{n-1}} - \frac{n^2(M_{II}-M_I)^2R^2}{2C_n^2 f(R)^2} - \frac{C_n^2 f(R)^2}{8n^2R^{2n}} \tag{25}$$

where $C_n \equiv \kappa\mu/S_n$.

Supposing that the function $n(L)$ has compact support, then we have $\alpha_\infty = 0$, and the asymptotic behavior of $V(R)$ would be

$$R \to 0 \qquad V(R) \to -\frac{C_n^2 N^2 \langle L\rangle^2}{8n^2R^{2n}} \tag{26}$$

$$R \to \infty \qquad V(R) \to \frac{C_n^2N^2 - n^2(M_{II}-M_I)^2}{2C_n^2N^2} \tag{27}$$

where $\langle L\rangle$ is the mean angular momentum modulus. We see that *the shell can have unbounded motion if and only if* $C_nN < n(M_{II} - M_I)$. Considering derivatives of (25), it can be shown in particular that *we can have oscillating or static solutions only if* $n = 2$, which is the case analyzed in [9]. It can also be shown that (14) implies that region II must be an *exterior* Schwarzschild region for both the unbounded and the oscillating solutions.

IV. STABILITY OF A TWO-COMPONENT HYPERSURFACE AGAINST SEPARATION OF THE CONSTITUENTS

We now consider a shell composed of two non-interacting matter fields. We impose that both matter fields are spherically symmetric, so their flow lines go with the comoving observers within the shell and they can be characterized as n-dimensional perfect fluids. The idea is to consider an infinitesimal separation of the components at a given proper time τ_0 and calculate the relative acceleration of the separating shells. The resulting space-time, once both components are infinitesimally separated, can be determined merely by imposing continuity of $R_0 \equiv R(\tau_0)$ and its first derivative, which is equivalent to impose smoothness (each "branch" should be an embedded submanifold) in the geometry of each "splitting" component, and also equivalent to impose continuity of normal vectors at the separation point.

The stress-energy tensor takes the form,

$$S^{ij} = (p_1 + p_2)h^{ij} + (\rho_1 + \rho_2 + p_1 + p_2)u^i u^j \tag{28}$$

Conservation of the source holds separately for each component,

$$\frac{d\rho_i}{dR} + \frac{n(\rho_i + p_i)}{R} = 0 \tag{29}$$

so we have in principle functions $\rho_i(R)$ and $p_i(R)$ for each component which are characteristic of the matter model. We should solve the equation of motion for the composed stress-energy tensor (28) and then for each component separately, taking into account the geometry that the new configuration would have, as illustrated in Fig.1. *If the resulting relative acceleration turns out to have the same sign than the relative displacement, then the shell is unstable, otherwise it should be considered as stable.*

In this Section we prove that certain families of shells, that are solutions of Einstein equations, are unstable against this kind of separation, while other families are stable. What will remain to be proven is whether there are solutions that are initially stable and evolve as a single shell up to a point where they *become* unstable and consequently could *undergo* splitting. In the next Section we prove by giving examples that such situations are possible, and splitting shells that are formally solutions of Einstein equations can be constructed.

The keys to analyze the relative acceleration of the resulting single-component shells are equation (15) and Birkhoff's theorem. The space-time region that would appear between the splitting shells is characterized by a mass parameter M_{int}. We label with the index 2 (ρ_2,p_2) the splitting shell that moves *into* region II. The equation of motion for the shell labeled with the index 1 is obtained by replacing $(M_I, M_{II}, \rho(R))$ in (13) with $(M_I, M_{int}, \rho_1(R))$; and for the shell labeled with the index 2 by replacing those quantities with $(M_{int}, M_{II}, \rho_2(R))$. Continuity of $\dot{R}(\tau_0)$ determines M_{int} as a function of (R_0, M_I, M_{II}) and the matter model parameters.

Looking at (11), it can be seen that positivity of ρ_1 and ρ_2 implies that M_{int} should be greater than at least one of the given mass parameters. In particular, it implies that if region II is *exterior*, then $M_I < M_{int} < M_{II}$, and $\nabla_a r$ in the intermediate region should point to region II (we label this possibility for the orientations with the A letter). In the same way, if region II is *interior*, then $M_{int} > M_{II} > M_I$, and both orientations for $\nabla_a r$ in the intermediate region are possible (we call B the case where $\nabla_a r$ points to II, and C the case where it points to I). In the latter case, in order to characterize the intermediate space-time, we must resort to the consistency condition (14) applied to one of the resulting shells. It can be shown that in any case the consistency conditions (14) applied for each splitting shell are indeed consistent with each other (in a given splitting scenario, M_{int} and $\nabla_a r$ in the intermediate region are always well-defined).



FIG. 1: Schematic representation of a *splitting* into two parts. The arrows inside the boxes represent the possible orientations of $\nabla^a r$ in each region. We distinguish the three possible sets of orientations (which we call A, B and C) compatible with the positivity of ρ_1 and ρ_2.

As we said, we can distinguish three possible splitting scenarios, depending on the character of region II and the region that would form between the separating components, as illustrated in Fig.1. We want to compute for each situation the relative acceleration of the shells, and to achieve this we propose using (15) adapted to each shell as follows,

$$\begin{aligned}
\ddot{R}_1(\tau_0) &= -\frac{n-1}{2R_0}\left[1+\dot{R}^2(\tau_0) - sign\left(\frac{\partial r}{\partial \eta_1}|_{int}\right)\left(\frac{2n}{n-1}\alpha_1+1\right)\sqrt{1+\dot{R}^2(\tau_0)-\frac{2M_I}{R_0^{n-1}}}\sqrt{1+\dot{R}^2(\tau_0)-\frac{2M_{int}}{R_0^{n-1}}}\right]\\
\ddot{R}_2(\tau_0) &= -\frac{n-1}{2R_0}\left[1+\dot{R}^2(\tau_0) - sign\left(\frac{\partial r}{\partial \eta_2}|_{II}\right) sign\left(\frac{\partial r}{\partial \eta_2}|_{int}\right)\left(\frac{2n}{n-1}\alpha_2+1\right)\sqrt{1+\dot{R}^2(\tau_0)-\frac{2M_{int}}{R_0^{n-1}}}\sqrt{1+\dot{R}^2(\tau_0)-\frac{2M_{II}}{R_0^{n-1}}}\right]
\end{aligned} \tag{30}$$

where $\alpha_i \equiv p_i/\rho_i$. We call the proper time coordinate of the original shell τ and that of the resulting shells τ_1 and τ_2, and make them coincide at the splitting point $\tau = \tau_1 = \tau_2 = \tau_0$.

These derivatives are with respect to different time coordinates, so in order to make a comparison we should rewrite them in terms of a single time. A natural choice in a splitting scenario would be the standard time coordinate of the Schwarzschild intermediate region t_{int}. To compute those derivatives we consider the following. For a general Schwarzschild space-time that has a shell as a boundary, and whose related variables and coefficients we note by a tilde, we can write,

$$\left(\frac{d\tau}{d\tilde{t}}\right)^2 = \frac{\tilde{F}^2}{\tilde{F}+\dot{R}^2} \tag{31}$$

where τ is the proper time coordinate of the boundary surface, R its radius, and $\tilde{F} \equiv 1-2\tilde{M}/R^{n-1}$. This expression implies that in a splitting scenario we have $d\tau_1/dt_{int} = d\tau_2/dt_{int}$ at τ_0. In this way, at a given proper time τ_0 we can write,

$$\frac{d^2R(\tau_0)}{d\tilde{t}^2} = \frac{1}{\tilde{F}}\left(\frac{d\tau}{d\tilde{t}}\right)^4 \ddot{R}(\tau_0) + \frac{\tilde{F}'}{2\tilde{F}}\left(\tilde{F}-\left(\frac{d\tau}{d\tilde{t}}\right)^2\right)\left(2\tilde{F}-\left(\frac{d\tau}{d\tilde{t}}\right)^2\right) \tag{32}$$

which in the splitting scenario implies

$$\left(\frac{d^2R_2}{dt_{int}^2}-\frac{d^2R_1}{dt_{int}^2}\right)|_{\tau=\tau_0}=\frac{F_{int}(R_0)^3}{(F_{int}(R_0)+\dot{R}(\tau_0)^2)^2}\left(\frac{d^2R_2}{d\tau_2^2}-\frac{d^2R_1}{d\tau_1^2}\right)|_{\tau=\tau_0}. \tag{33}$$

We now can see that, as a consequence of the continuity of $R(\tau)$ and it first derivative, the stability condition turns out to be simply

$$sign\left(\frac{\partial r}{\partial\eta_1}|_{int}\right)(\ddot{R}_2(\tau_0)-\ddot{R}_1(\tau_0))<0. \tag{34}$$

Taking into account (30) and (11) for each splitting shell, this inequality takes the form

$$\beta\equiv\kappa^2(\rho_1+\rho_2)^2R^{n+1}(n(\alpha_1+\alpha_2+1)-1)-2n^3(M_{II}-M_I)(\alpha_2-\alpha_1)>0 \tag{35}$$

which is independent of the signs of $\partial r/\partial\eta|_{int}$ and $\partial r/\partial\eta|_{II}$. It is remarkable that *in all the three cases A, B and C, the stability condition takes the form (35).* As a first result, we note that this condition implies that *if both fluids are identical ($\alpha_2=\alpha_1=\alpha$) and $\alpha>-(n-1)/2n$, then the shell is always stable.* A somehow unexpected result is that if we have two identical fluids with $\alpha<-(n-1)/2n$, then the system is unstable. That would mean that if the matter content of a shell satisfies the latter inequality and it makes sense to separate the stress-energy tensor in two equal-alpha parts, then such a shell would always be unstable. We also note that if both $\alpha_i\geq 0$ (which is the case for matter made of particles), then there is instability only if α_2 is large enough in comparison to α_1. In the next subsections we particularize the stability analysis for some specific matter models.

Fluids with equation of state $p_i=\omega_i\rho_i$

We propose two fluids with equation of state $p_i=\omega_i\rho_i$, where $\omega_2>\omega_1$. In this case (35) takes the form

$$(C_1R^{-n(1+\omega_1)}+C_2R^{-n(1+\omega_2)})^2R^{n+1}(n(\omega_1+\omega_2+1)-1)-2n^3(M_{II}-M_I)(\omega_2-\omega_1)>0. \tag{36}$$

Provided $\omega_1>-(n-1)/2n$, then *the shell always become unstable for R large enough.*

As we have seen in the previous Section, at large R the ω_1 fluid dominates $\rho(R)$, and the asymptotic behavior is determined by $\alpha_\infty=\omega_1$, so there are solutions with R arbitrarily large provided $\omega_1\geq 0$. We are proving in this way that there exist shells so that, while being solutions of Einstein equations, are unstable and therefore, as we will argue later on, unphysical.

Vlasov matter

In the case of Vlasov matter, what characterizes the matter distribution is the function $n(L)$. We may define non-interacting components merely by separating the particles into two sets that satisfy spherical symmetry on their own. For each set we would have an angular momentum distribution function $n_i(L)$ so that $n_1(L)+n_2(L)=n(L)$. We may define for each set,

$$\alpha_i=\frac{1}{\int n_i(L)\sqrt{R^2+L^2}dL}\int\frac{n_i(L)L^2}{n\sqrt{R^2+L^2}} \tag{37}$$

where we can see that $0\leq\alpha_i\leq 1/n$. We commented that instability appears whenever $\alpha_2-\alpha_1$ is large enough. That could be achieved if one of the sets, say $n_2(L)$, comprises all the particles with L greater than a certain value, as illustrated by the following case.

In particular, we analyze perhaps the simplest case that could display this kind of instability, a two-component shell where $n_i(L) = N_i\delta(L - L_i)$. The condition (35) reads,

$$C_n^2 f(R)^2[(n+1)L_1^2L_2^2 + nR^2(L_1^2 + L_2^2) + (n-1)R^4] - 2n^2R^{n+3}(M_{II} - M_I)(L_2^2 - L_1^2) > 0 \quad (38)$$

where we can see that for R large enough ($N = N_1 + N_2$):

- $n = 2$, the shell is stable,
- $n = 3$, the shell is stable if $\kappa^2\mu^2N^2 - 36\pi^4(M_{II} - M_I)(L_2^2 - L_1^2) \geq 0$ and unstable otherwise,
- $n > 3$, the shell is unstable.

As we discussed in the previous section, there are shells that go to infinity provided $n(M_{II} - M_I) > C_nN$. So a shell satisfying this last inequality and $n > 3$, or $n = 3$ and $\kappa^2\mu^2N^2 < 36\pi^4(M_{II} - M_I)(L_2^2 - L_1^2)$, satisfies Einstein equations while being unstable at R large enough. In the next Section we give explicit examples where the shells are initially stable, become unstable at a later time, and where splitting solutions can be constructed.

V. SPLITTING SHELLS

In this Section we construct splitting solutions where the splitting takes place at a critical point of stability condition (35) (when the equality holds). These shells *become* unstable at that time, and a *soft* splitting solution may be built, that is, a splitting solution where $\ddot{R}$ is also continuous (which is what defines the critical condition).

Fluids with equation of state $p_i = \omega_i\rho_i$

As a first example we consider a shell made of two non-interacting fluids, whose parameters are as follows.

- **Example 1**: $n = 2$, $\kappa = 8\pi$, $\omega_1 = 0$, $\omega_2 = 0.001$, $C_1 = 0.01$, $C_2 = 0.02$, $M_I = 1$, $M_{II} = 2$

We illustrate in Fig.2 the velocity of the shell and the stability condition as functions of R. There are no prohibited regions for the motion of this shell, so if it is initially expanding then it will continue to expand forever (provided the system evolves as a single shell during its entire evolution). We can see that at $R_c \approx 35,27$ the shell *becomes* unstable. If we consider an initial radius $R_0 < R_c$, and say that the shell is initially expanding, then at $R = R_c$ we may construct a splitting of the type illustrated in Fig.1, where the M_{int} parameter and the orientation of the r coordinate would be fixed by the continuity of $\dot{R}$ and the conditions (14) applied to each spitting shell respectively. As region II must be exterior, and the energy densities of each component must be positive, the r coordinate turns out to be a global coordinate (type C splitting).

Vlasov matter

As a second example we consider a shell made of Vlasov matter, such that the particles only have two possible values for the modulus of their angular momentum ($n(L) = N_1\delta(L - L_1) + N_2\delta(L - L_2)$). An example of this kind was already constructed in [9] for a 4-dimensional space-time ($n = 2$). In this paper we build a 5-dimensional example (like the previous one) whose parameters satisfy both $n(M_{II} - M_I) > C_nN$ and $\kappa^2\mu^2N^2 < 36\pi^4(M_{II} - M_I)(L_2^2 - L_1^2)$.



FIG. 2: The solid curve represents $\dot{R}^2$ as a function of R for a shell made of two non-interacting fluids with parameters $n = 2$, $\kappa = 8\pi$, $\omega_1 = 0$, $\omega_2 = 0.001$, $C_1 = 0.01$, $C_2 = 0.02$, $M_I = 1$ and $M_{II} = 2$. The dotted curve represents 1000β. The critical point is located at $R \approx 35,27$; for R greater than that the shell results unstable.

- **Example 2:** $n = 3$, $\kappa = 1$, $M_I = 1$, $M_{II} = 3$, $\mu N_1 = 30$, $\mu N_2 = 23$, $L_1 = 2$, $L_2 = 2.1$

We illustrate the effective potential $V_2(R)$ for this example in Fig.3. There are no prohibited regions for this shell. We propose a separation between the particles with L_1 and the particles with L_2. At $R_c \approx 27,98$ there is a critical point of the stability condition (35), and for R greater than that the shell is unstable. As in the previous case, we propose a soft splitting at the critical point, provided $R_0 < R_c$ and that the shell is initially expanding, and obtain M_{int} from the continuity of $\dot{R}$. In this case it also turns out that the r coordinate is a global coordinate. In Fig.3 we illustrate the differences between the effective potentials associated with each splitting component and the one associated with the original shell. As Newtonian intuition suggest, we have that the shell composed of particles with angular momentum L_2 begins to expand faster than the rate at which the original shell would expand, while the shell with L_1 begins to expand slower than that rate.

VI. THREE-COMPONENT SHELL IN A Z_2 SYMMETRIC SCENARIO

In the context of brane cosmology, a common simplifying assumption coming from string theory is the existence of Z_2 symmetry for the bulk space-time, such that the brane-world is the set of fixed points of this symmetry [3]. In this case, (14) confirms that region II must have the same character than region I, they both must be *interior* Schwarzschild regions with identical mass parameters.

We consider a different kind of splitting in the Z_2 symmetric scenario, as illustrated if Fig.5. To preserve the symmetry we propose a fragmentation into three parts, one of which remains at the symmetry center, while the other two, being mirror images of one another, move away from that symmetry center. This is a three-component (non-interacting) shell splitting, where two of them are identical. We can consider the "outgoing" shells as two non-interacting identical parts of a single matter-energy component, whenever such interpretation makes sense. Analogously to Section IV, the key to analyze this kind of instability is to use (15) in order to obtain the relative acceleration of the resulting shells. We call M the mass parameter of the Schwarzschild regions



FIG. 3: $V_2(R)$ for $n = 3$, $\kappa = 1$, $M_I = 1$, $M_{II} = 3$, $\mu N_1 = 30$, $\mu N_2 = 23$, $L_1 = 2$, $L_2 = 2.1$. There are no prohibited regions, and the curve is already very close to its asymptote at $R \approx 20$.



FIG. 4: The dotted curve depicts $V_2(R; \rho_1, M_I, M_{int}) - V_2(R; \rho, M_I, M_{II})$, while the dashed curve depicts $V_2(R; \rho, M_I, M_{II}) - V_2(R; \rho_2, M_{int}, M_{II})$. A solution representing a *splitting* like that illustrated in Fig.1 can be constructed, where the separation takes place at $R \approx 27,98$. After the separation, the shell with $L_2 = 2.1$ would expand faster than the shell with $L_1 = 2$.

before the splitting, and M' the mass parameter that would correspond to the region between the symmetry center and one of the "outgoing" shells. We now define ρ_0 and p_0 as the stress-energy tensor components of the central splitting shell, and denote the corresponding components for the outgoing shells by $\tilde{\rho}/2$ and $\tilde{p}/2$, so we can write $\rho = \rho_0 + \tilde{\rho}$ and $p = p_0 + \tilde{p}$. The equation of motion of the resulting central shell is obtained by replacing $(M_I, M_{II}, \rho(R))$ with $(M', M', \rho_0(R))$ in (13), and for the outgoing shells by replacing those parameters with $(M', M, \tilde{\rho}(R)/2)$ in the same equation. Continuity of $\dot{R}(\tau_0)$ determines M' as a function of (M, R_0) and the matter models parameters, and it can be shown that M' is always well-defined. Positivity of each energy density component implies that $\nabla_a r$ always points to the central brane, which in turn implies $M' > M$.



FIG. 5: Schematic representation of a *splitting* into three parts that preserves Z_2 symmetry. The arrows in each empty region represent the orientations of $\nabla_a r$.

We have for each shell,

$$\begin{aligned}\ddot{\tilde{R}}(\tau_0) &= -\frac{n-1}{2R_0}\left[1+\dot{R}^2(\tau_0)-\left(\frac{2n}{n-1}\tilde{\alpha}+1\right)\sqrt{1+\dot{R}^2(\tau_0)-\frac{2M}{R_0^{n-1}}}\sqrt{1+\dot{R}^2(\tau_0)-\frac{2M'}{R_0^{n-1}}}\right] \\ \ddot{R}_0(\tau_0) &= -\frac{n(\alpha_0+1)-1}{R_0}\left(1+\dot{R}^2(\tau_0)-\frac{2M'}{R_0^{n-1}}\right)-\frac{(n-1)M'}{R_0^n} \end{aligned} \tag{39}$$

Analogously to Section IV, the condition for having stability would be simply

$$\ddot{\tilde{R}}(\tau_0)-\ddot{R}_0(\tau_0)>0. \tag{40}$$

After manipulating (39) and taking into account that $\rho>0$ and $\rho_0>0$, we finally get

$$(2n\tilde{\alpha}+n-1)(\rho_0+\tilde{\rho})+(2n\alpha_0+n-1)\rho_0>0. \tag{41}$$

This inequality can also be written as,

$$2n(p_0+\rho_0)+2(n\tilde{\alpha}-1)\rho_0+(2n\tilde{\alpha}+n-1)\tilde{\rho}>0 \tag{42}$$

where it is clear that *if $\tilde{\alpha}>1/n$, then the shell is always stable.* We now suppose that on the original shell there is brane tension plus matter satisfying the dominant energy condition, so we may write $\rho_0=b+\rho_m$, $p_0=-b+p_m$, which illustrates the fact that the resulting central shell must also have brane tension[3]. In this case, we can rewrite the inequality (41) as follows

$$2(n(\alpha_m+\tilde{\alpha}+1)-1)\rho_m+2(n\tilde{\alpha}-1)b+(2n\tilde{\alpha}+n-1)\tilde{\rho}>0. \tag{43}$$

We can see that if $0\leq\tilde{\alpha}\leq 1/n$, and $0\leq\tilde{\alpha_m}\leq 1/n$, then *the shell is unstable for $b>>\rho_m$ and $b>>\tilde{\rho}$.* Hence, if the shell is composed of non-interacting particles (Vlasov matter) plus some other matter field (ρ_m, p_m) and a dominating brane tension, then *it is typically unstable against separation of components*. In that case, $(\tilde{\rho}, \tilde{p})$ would be determined by (23)), and we may separate the particle ensemble in two identical parts $(n_1(L) = n_2(L) = n(L)/2)$ in order to construct a splitting of this kind. In typical brane-world models, b turns out to be much greater than any other energy density at least since nucleosynthesis, so it is relevant to ask whether this kind of instability also appears in brane cosmology scenarios, which is the subject of the next Section.

---

[3] If we consider that the brane tension component is indivisible, it must remain at the symmetry center.

VII. SPLITTING BRANE-WORLDS

In this Section we extend the instability analysis that we made in previous Sections to actual brane-world scenarios. We consider SMS type brane-worlds [10] in which the five-dimensional bulk space-time has a spacial 4-slicing in which every slice possesses a kind of 3-dimensional isotropy: spherical, planar or hyperbolic symmetry. For such space-time, every timelike surface defined in the quotient manifold of the group orbits is a FRW submanifold. Bowcock, Charmousis and Gregory [11] have shown that this kind of manifold, provided they are solutions of vacuum Einstein equations with a cosmological constant, are always 5-dimensional Schwarzschild-Anti de Sitter space-times. This is the main property that allows us to extend the analysis that we have done for spherically symmetric space-times, as it plays the role that Birkhoff's theorem played in Sections IV and VI. In this way, every open vacuum region in the bulk must be Schwarzschild-AdS, which is a static space-time characterized by two parameters: a mass parameter and the cosmological constant Λ_5. The metric in standard static coordinates for this kind of space-time reads,

$$ds^2 = -F(r)dt^2 + F(r)^{-1}dr^2 + r^2\left[\frac{d\chi^2}{1-k\chi^2} + \chi^2 d\Omega^2\right] \tag{44}$$

where $F(r) = k - 2M/r^2 + r^2/\ell^2$, k is the curvature index $(1, 0, -1)$ and $\ell^2 \equiv -6/\Lambda_5$.

In this context, the brane-world is a FRW timelike boundary layer separating two different Schwarzschild-Anti de Sitter regions (I and II). We consider for simplicity the same cosmological constant at both sides, but eventually different mass parameters. Like in the previous Sections, this singular surface can be described by a timelike curve in a quotient manifold, so a proper time coordinate τ can be naturally defined, and the extrinsic curvature in $(\tau, \chi, \theta, \phi)$ coordinates it is also expressed by (6). This coordinate system on the brane-world submanifold are precisely the standard coordinates for FRW space-times, where the scale factor is simply $a(\tau) = r(\tau, \eta = 0) \equiv R(\tau)$ (η is the gaussian normal coordinate). As in the previous analysis, the matter-energy content of the shell is described by two functions ρ and p, as expressed by (8). Then, Israel-Darmois junction conditions applied to this situation could also be written as (11) and (12). Analogously, we suppose that at least one of the vacuum bulk regions is "interior", which means that it can be described by an inequality $r < R(t)$ in the static coordinate, and which is also a consequence of imposing $\rho > 0$ matter fields on the brane[4].

In this way, as follows from (11), there is an equation of motion for the brane like that of a particle with unit mass and zero energy in a potential given by,

$$V(R) = \frac{k}{2} - \frac{M_I + M_{II}}{2R^2} + \frac{R^2}{2\ell^2} - \frac{9(M_{II} - M_I)^2}{2\kappa^2\rho^2 R^6} - \frac{\kappa^2\rho^2 R^2}{72} \tag{45}$$

and that equation of motion can also be written as follows

$$H^2 \equiv \left(\frac{\dot{R}}{R}\right)^2 = \frac{\kappa^2\rho^2}{36} + \frac{9(M_{II} - M_I)^2}{\kappa^2\rho^2 R^8} + \frac{M_I + M_{II}}{R^4} - \frac{1}{\ell^2} - \frac{k}{R^2}. \tag{46}$$

---

[4] Provided $R > 2max\{M_I, M_{II}\}$.

In order to recover the classical Friedmann equations as a low energy limit, brane tension is needed $(\rho = \rho_m + b, p = -b + p_m)$. In that case, (46) can be written [13],

$$H^2 = \frac{\mathcal{C}}{(\rho_m/b+1)^2 R^8} + \frac{8\pi G}{3}\rho_m\left(1+\frac{\rho_m}{2b}\right) - \frac{k}{R^2} + \frac{M_I + M_{II}}{R^4} + \frac{\Lambda_4}{3} \tag{47}$$

where $G \equiv \kappa^2 b/48\pi$, $\Lambda_4 \equiv (\kappa^2 b^2 + 6\Lambda_5)/12$, and $\mathcal{C} \equiv 3(M_{II} - M_I)^2/16\pi G b$. We notice that there is a "dark radiation" term and a term related to the "asymmetry" of the bulk space-time which may cause an inflation-like stage at early $(\rho_m > b)$ times[5]. Because we consider a uniform bulk cosmological constant, (14) would determine whether a given solution of the equation of motion is actually a solution of the junction conditions as in the spherically symmetric case. We can see that in the case of Z_2 symmetry centered on the brane, the consistency condition always holds (both empty bulk regions are identical, hence *interior*); while for general non-Z_2 symmetric brane-world scenarios, (14) would typically imply that both bulk vacuum regions must also be *interior*, because of the observational constraints on dark radiation (which are constraints on the $(M_I + M_{II})/R^4$ term).

A. Instability against separation of components in non-Z_2-symmetric scenarios

We now propose a brane-world made of two non-interacting homogeneous and isotropic matter-energy components, so we may write $\rho = \rho_1 + \rho_2$ and $p = p_1 + p_2$. One of this components could always be the brane tension $\rho_i = -p_i = b$. In this subsection we consider a splitting like that of Fig.1, and impose continuity of the normal vectors at the separation point. We assume that the whole space-time possesses the symmetries we imposed above, so the intermediate region has to be Schwarzschild-AdS, with some (M, Λ_5) parameters. In this paper we are not considering the possibility of having different cosmological constants on each open vacuum region, so the geometry of the intermediate region would be determined by the mass parameter M_{int} and the orientation of ∇r on it. As in Section IV, both M_{int} and ∇r are uniquely determined by the junction and consistency conditions applied to each splitting component.

The junction condition on the brane (12) implies,

$$\ddot{R} = -\frac{1}{R}\left[k + \frac{2R^2}{\ell^2} + \dot{R}^2 \mp (3\alpha+1)\sqrt{\dot{R}^2 + F_I}\sqrt{\dot{R}^2 + F_{II}}\right] \tag{48}$$

which can be applied to each splitting component in order to determine the relative acceleration of the separating shells. Taking the same steps as in Section IV, the stability condition turns out to be precisely (34), which in this context can be written,

$$54(M_{II} - M_I)(\alpha_2 - \alpha_1) - \kappa^2\rho^2 R^4(3(\alpha_1+\alpha_2)+2) < 0 \tag{49}$$

regardless of the relative orientation of ∇r in the intermediate region. Now, if we consider that the brane tension stress-energy should be conserved, one of the two splitting shells should have brane tension. In this way, we may write $\alpha_1 = (-b + p_{1m})/(b + \rho_{1m})$ which in the low-energy limit simply becomes $\alpha_1 \approx -1$, so the stability condition can be written

$$\kappa^2 b^2 R^4(3\alpha_2 - 1) - 54(M_{II} - M_I)(\alpha_2 + 1) > 0. \tag{50}$$

Because there are very tight observational bounds on dark radiation, this stability condition for "realistic" brane-world scenarios turns out to be merely $\alpha_2 > 1/3$, which is a condition that matter made of particles does not satisfy. We may consider (ρ_2, p_2) as the matter-energy content of the universe, or some non-interacting part of it (like the dark matter component to a good approximation), which is thought to have always satisfied $\alpha < 1/3$, and indeed it practically satisfies $\alpha \approx 0$ in the matter-dominated era. We then conclude that *non-Z_2-symmetric brane-world scenarios are typically unstable against separation of non-interacting components.*

---

[5] Which nevertheless can not account actual inflation, as explained in [14]

Although the name of this subsection makes reference to non-Z_2-symmetric scenarios, this kind of instability would also hold for an *initially* Z_2 symmetric bulk space-time. Condition (50) makes perfect sense if $M_I = M_{II}$, which is equivalent in this context to having Z_2 symmetry. In that case, a two-component splitting would necessarily break the symmetry, so if we have a reason to impose it, as it may happen in brane-world models, such a splitting scenario should not be taken into account.

B. Instability against separation of a three-component brane-world in a Z_2-symmetric scenario

Nevertheless, as we did in Section VI, we may propose a splitting of a shell made of three non-interacting components that preserves Z_2 symmetry, as illustrated in Fig.5. The mass parameter of the intermediate region M' would be determined by the junction conditions and the continuity of normal vectors at the separation point. By means of an analysis completely analogous to that of Section VI, the stability condition turns out to be,

$$(3\alpha_0 + 1)\rho_0 + (3\tilde{\alpha} + 1)(\tilde{\rho} + \rho_0) > 0 \tag{51}$$

where the 0 index denotes the component that remains to be at the symmetry center, while the tilde denotes quantities that correspond to the matter conforming the two symmetrically "outgoing" shells. The brane tension component must remain within the central brane, so we can write $\rho_0 = b + \rho_{0m}$ and $p_0 = -b + p_{0m}$. In the low energy limit, the stability condition for this splitting scenario turns out to be $\tilde{\alpha} > 1/3$, which is the same condition than that for a two-component splitting. As we argued, if it makes sense to separate some non-interacting component of the universe into two non-interacting and identical parts, something that in particular always makes sense for dark matter, then splitting branes can be constructed, that are solutions of Einstein equations suggesting a particular kind of instability for brane-worlds, as we will discuss later on.

C. A "realistic" example

In this subsection we want to illustrate with an example how to construct a splitting solution in this context. We consider a SMS brane-world universe whose parameters are in agreement with recent observations, interpreted in the *concordance model* framework. We consider a brane-world universe whose matter content resembles that of the $\Lambda - CDM$ universe, so we may write,

$$\rho_m = \rho_c(\Omega_c + \Omega_b)(1+z)^3 + \epsilon_0^{\gamma}(1+z)^4 \tag{52}$$

where $\rho_c \equiv 3H_0^2/8\pi G$ is the critical density at present cosmological time, Ω_c and Ω_b are the relative proportions of dark matter and baryonic matter respectively at present with respect to the critical density, and ϵ_0^{γ} is the present radiation energy density. For simplicity, we consider the simplest choice of parameters ($k = M = 0$) for both empty bulk regions, which particularly implies non-existence of "dark radiation" and Z_2 symmetry (each bulk region is an identical piece of AdS_5).

The Friedmann equation for such a universe reads,

$$H^2 = H_0^2\left[\left((\Omega_c+\Omega_b)(1+z)^3+\frac{\epsilon_0^\gamma}{\rho_c}(1+z)^4\right)\left(1+\frac{\epsilon_0^\gamma(1+z)^4}{2b}\right)+\Omega_\Lambda\right] \tag{53}$$

where $\Omega_\Lambda \equiv \Lambda_4/(3H_0^2)$, and we have ignored every term other than the radiation term in the quadratic in ρ_m part of this expression. For small z, this equation takes the standard form,

$$H^2 = H_0^2((\Omega_c+\Omega_b)(1+z)^3+\Omega_\Lambda) \tag{54}$$

which makes explicit the relation $\Omega_c+\Omega_b+\Omega_\Lambda = 1$.

We propose a splitting in three parts like that of Section VI, where the $(\tilde{\rho},\tilde{p})$ *component is the dark matter component* $(\tilde{\rho}=\rho_c\Omega_c(1+z)^3, \tilde{p}=0)$ *and the splitting takes place at present cosmological time* $(z=0)$*.* It makes sense to separate two identical dark matter components, which would constitute the two "outgoing" shells, as cold dark matter is supposed to be made of non-interacting particles, hence it can be modeled as Vlasov matter and in this case (at cosmological scales) simply as dust. We should emphasize however that we do not expect this kind of construction to have *a priori* any degree of correspondence with the actual universe., They are solutions of Einstein equations that correspond to initial value data that can be obtained from a family of brane-world models analyzed in the literature. As actual solutions, we may consider the phenomenology that one would expect for observers living in a splitting brane-world, just to explore the theoretical possibility. In any case, with this construction we provide another example of the *lack of uniqueness* in the evolution of initial data that corresponds to concentrated sources.

As we commented, brane-world models are typically unstable against separation of components. Replacing ρ with $\tilde{\rho}/2$ and setting $(M_I=0, M_{II}=M')$ in (46), the equation of motion for any of the "out-going" dust shells can be written as follows,

$$\tilde{H}^2 = \frac{8\pi G}{3}\tilde{\epsilon}_{dr}\left(1+\frac{2b\tilde{\epsilon}_{dr}}{\tilde{\rho}^2}\right)+\frac{\pi G}{3b}\tilde{\rho}^2+\frac{\Lambda_4-4\pi G b}{3} \tag{55}$$

where we replaced κ and Λ_5 as functions of G and b, and $8\pi G\tilde{\epsilon}_{dr}/3 \equiv M'/R^4$ is the "dark radiation" term associated with the mass parameter M' of the intermediate region. Taking into account the fact that $\tilde{\rho} << b$ at the separation moment, we can write

$$\tilde{H}^2 = H_0^2\left[\tilde{\Omega}_{dr}(1+\tilde{z})^4\left(1+\frac{2b\tilde{\Omega}_{dr}}{\rho_c\Omega_c^2(1+\tilde{z})^2}\right)+\Omega_\Lambda-\frac{b}{2\rho_c}\right] \tag{56}$$

where $\tilde{\Omega}_{dr} \equiv \tilde{\epsilon}_{dr}(R_0)/\rho_c$ and $\tilde{z} \equiv R_0/\tilde{R}-1$. In this way, all we need to determine the equation of motion of the resulting dust shells is the parameter $\tilde{\Omega}_{dr}$, which can be found by continuity of the normal vectors at the separation point, that is, by solving it from $H(z=0)=\tilde{H}(\tilde{z}=0)$. From (54) and (56) we get,

$$\tilde{\Omega}_{dr}^2 \approx \frac{\Omega_c^2}{4}\left(\frac{\rho_c}{b}(2\Omega_b+\Omega_c)+1\right) \tag{57}$$

so (56) can be written

$$\tilde{H}^2 = H_0^2\left[\left(\frac{b}{2\rho_c}+\frac{\Omega_c}{2}+\Omega_b\right)(1+\tilde{z})^2+\frac{\Omega_c}{2}(1+\tilde{z})^4+\Omega_\Lambda-\frac{b}{2\rho_c}\right]. \tag{58}$$

This expression can be simplified neglecting terms $(b >> \rho_c)$, so we can write,

$$\tilde{H}^2 = H_0^2\left[1+\frac{b}{\rho_c}\tilde{z}\left(1+\frac{\tilde{z}}{2}\right)\right] \tag{59}$$

where it can be easily seen that the dust shells almost immediately rebound ($\tilde{H} = 0$ at $\tilde{z} \approx -\rho_c/b$), which implies that the shells scale parameter begins to decrease until they collapse and form a pair of bulk black holes with mass parameter M'.

On the other hand, the equation of motion of the central shell after the splitting can be computed from (47) by writing $\rho_m = \Omega_b(1+z)^3$ and $M_{II} = M_I = M'$,

$$H^2 = H_0^2 \left[\Omega_b(1+z)^3 + \Omega_c(1+z)^4 + \Omega_\Lambda\right] \tag{60}$$

where the second term of the right hand side corresponds to the "dark radiation" term M'/R^4. In this way, the motion of the central shell is determined by (54) for $z > 0$ and by (60) for $z < 0$. As we can see, the time derivative of the Hubble parameter would be discontinuous for the central shell, and the dynamical effect of the "disappearance" of the dark matter component would be *as if the dark matter of the universe transformed into dark radiation*, as the initial ($z = 0$) effective energy density of dark radiation is precisely the dark matter density at the time of the splitting.

VIII. CONCLUDING REMARKS

As far as we know, this kind of phenomena in the context of brane-world cosmologies was not previously considered in the literature. Nevertheless, there might be reasons in the context of string theory to *a priori* preclude any chance of having matter moving off the brane, as one may consider branes as fundamental objects by themselves, which appear in the definition of the action that describes the theory one is seeking to build. We basically affirm that these systems, in the framework of General Relativity, are unstable, or at least they entail non-uniqueness in the evolution, and the existence of those features may be an argument against the plausibility of the brane-world models considered in this work, and of certain families of singular hypersurface models in general. We remark that the critical point for any of the stability conditions (50,51) takes place at very early stages of the universe, long before nucleosynthesis. In this way, the brane results unstable almost during its entire evolution, and becomes increasingly unstable as time passes. A "soft" splitting, like those illustrated in Section V, where $\ddot{R}$ is continuous, can only take place at the critical point of the stability condition, so it could only happen in the inflationary era (taking into account experimental bounds on the brane tension based on deviations of Newtonian gravitation at submillimeter scales [18]), or even before. We may speculate in further investigations the phenomenological consequences of having this kind of splitting in the context of the very early universe.

In any case, this problem should be related to the confinement of particles in brane-world models, as it happens in the case of Vlasov matter in 4-dimensional space-time [9]. In Section 5 of that paper, the stability against single particle evaporation was considered. It was then concluded that for two-component Vlasov shells (like the one described in the second example of Section V) *the stability condition for separation of components is always stronger than the stability condition for single particle evaporation.* We may conjecture that for Vlasov matter the confinement of particles is a necessary condition for stability against separation of components. Nevertheless, the particle evaporation analysis may be relevant on its own, as the concept of "particle" in the cosmological context may have more sense physically than the concept of "non-interacting-component". Particles in the universe, both if they are dark or baryonic matter, may follow geodesics of the induced metric to a good approximation (as long as collisions can be neglected). But, as Anderson and Tavakol analyzed in [16] and we pointed out in [9], *geodesics of the induced metric do not generally coincide with geodesics of the bulk metric.* Indeed, the geodesics of the bulk metric on the brane are not uniquely defined because of the discontinuity of the affine connection: for a given initial point on the singular surface and a given initial tangent velocity there might be two geodesics (each one going into opposed bulk regions), a single geodesic, or no geodesic at all. Gravitational confinement is related to the non-existence of bulk geodesic for freely falling observers on the brane, as it means that according to the connection of region II the geodesic goes into region I and viceversa[6]. *If there exist a bulk geodesic and there are no non-gravitational forces acting on that particle, then it should classically follow that geodesic.* As analyzed in [17], the confinement of freely-falling observers to domain walls in Z_2-symmetric models can be accomplished if and only if the matter content of the shell satisfy the strong energy condition, which is a condition that models with brane tension generally do not satisfy, so it is no surprising that these models turn out to be unstable against separation of components.

We could also interpret the instability result as an evidence for the existence of some mechanism that would "prevent" the splitting, that is some interaction among the constituents that may compensate the instability. We may take into account for example a scalar field on the bulk, which is a widely considered scenario as it may naturally exhibit an inflationary stage and it is easily incorporated by string theoretical considerations. Another possibility is to consider Lovelock gravity for the bulk. These are possible future research directions, and it may happen that some brane-world scenarios turn out to be stable in this sense, so that the stability against separation of constituents could work as an argument in favor of some of them. As a first step, as we suggested, we may look into brane-world models that naturally incorporate particle confinement.

Another interesting issue regarding these solutions is the *lack of uniqueness* in the evolution of boundary layers. We have seen different situations involving singular hypersurfaces where a splitting solution can be constructed. But the splitting prescription is something that we can not see by looking at initial data in Cauchy surfaces that intersect the singular surface before the separation point. For every splitting solution that we constructed in this work, there is a corresponding non-splitting solution where the shell moves according to a equation of motion, and it may collapse, oscillate or expand always as a whole. That initial data is exactly the same for the splitting solution as for the corresponding non-splitting solution, that is, we do not have uniqueness in the evolution. Furthermore, if we consider particle evaporation we could also built solutions where an *atmosphere* forms, creating even more possible evolutions. This result may not be surprising taking into account the distributional data that a thin shell represents. In other contexts involving hyperbolic PDE, like ideal fluids in Minkowski space-time, surface of discontinuity may form and evolve (shock-waves) and their evolution according to the original PDE may not be unique [19]. There is an external principle (an entropy law) that recovers uniqueness. In this context we may have the same situation, as one may identify the "physical" evolution among the mathematical possibilities by means of some principle arising from the specific physics of the matter-energy content of the thin shell. After all, the information about the splitting may be exclusively contained in the degrees of freedom associated with the constituents of the singular surface. A specific treatment of these issues should be considered in a separate paper.

On the other hand, looking at the results in [20], where we shown that certain singular surfaces are indeed thin shell limits of thick configurations, and that all those singular surfaces are stable against separation of constituents, we make the following conjecture: *unstable shells are not thin shell limits of thick configurations.* If we consider thin shells as idealizations of ultimately thick configurations, those unstable solutions could be interpreted as bad idealizations of thick configurations, and hence they can only be taken into account if we consider the system to be *fundamentally thin*, that is, if there is some fundamental principle that simply precludes the possibility of having matter-energy moving off the surface. An unstable thin shell may be considered as a *spurious solution* arising from the lost of information associated with the fact that we neglected a spacial dimension. A general demonstration of this conjecture constitutes a possible future research direction, and it should be deeply related to the general initial value problem for concentrated sources. In any case, the stability-against-separation analysis that we exposed in this work can be regarded as a novel analysis to be taken into account in order to constraint the plausibility of models involving singular surfaces.

---

[6] The case where the bulk geodesics are contained within the brane are typically of zero-measure.

IX. ACKNOWLEDGMENTS

This work was partly supported by CONICET. I wish to thank Reinaldo Gleiser for inspiration and insightful ideas, Robert Geroch for illuminating discussions and advice, and Oscar Reula for useful discussions regarding the initial value problem for shock waves.

[1] Israel W, *Nuov. Cim.* B **44**, 1 (1966).
[2] Randall L, Sundrum R, *Phys. Rev. Lett.* **83**, 4690-4693 (1999).
[3] Randall L, Sundrum R, *Phys. Rev. Lett.* **83**, 3370-3373 (1999).
[4] Maartens R, Koyama K, “Brane-World Gravity” *Living Rev. Rel.* **13**, 5 (2010) *http://www.livingreviews.org/lrr-2010-5.*
[5] Barrabes C, Israel W, Letelier P S, *Phys. Lett. A* **160**, 41-44 (1991).
[6] Yangurazova L R, Bisnovatyi-Kogan G S, *Astroph. and Space Sci.* **100**, 319-328 (1984).
[7] Berezin V A, Kuzmin V A, Tkavech I I, *Phys. Rev. D* **36**, 10 2919 (1987)
Ansoldi S, Guendelman E, Shilon S, *Proceedings of BH2: Dynamics and Thermodynamics of Black Holes and Naked Singularities* - Politecnico di Milano, Italy (2007).
[8] Lukas A, Ovrut B A, Stelle K S, Waldram D, *Phys. Rev. D* **59**, 086001 (1999).
[9] Gleiser R J, Ramirez M A, *Class. Quant. Grav.* **26**, 045006 (2009).
[10] Shiromizu T, Maeda K, Sasaki M, *Phys. Rev. D* **62** 024012, 1-6 (2000).
[11] Bowcock P, Charmousis C, Gregory R, *Class. Quant. Grav.* **17**, 4745-4763 (2000).
[12] Fayos F, Senovilla J M M, Torres R, *Phys. Rev. D* **54**, 4862-4872 (1996).
[13] Stoyca H, Tye S-H H, Wasserman I, *Phys Lett. B* **482**, 205-212 (2000).
[14] Davis A-C, Davis S C, Perkins W B, Vernon I R, *Phys. Lett. B* **504**, 254 (2001).
[15] Andréasson H, “The Einstein-Vlasov system/ Kinetic theory” *Living Rev. Rel.* **14**, 4 (2011).
[16] Anderson E, Tavakol R, *JCAP* **0510**, 017 (2005).
[17] Seahra S S, Wesson P S, *Class. Quant. Grav.* **20**, 1321-1340 (2003).
[18] Long J C, Chan H W, Churnside A B, Gulbis E A, Varney M C M, Price J C, *Nature* **421**, 922-925 (2003).
[19] Christodoulou D, “The formation of shocks in 3-dimensional fluids”, *European Mathematical Society* (2007).
[20] Gleiser R J, Ramirez M A, *Class. Quant. Grav.* **27**, 065008 (2010).